

Table 3: COVID-19 vaccination schedule for people with moderate or severe immunocompromise*

Primary vaccination	Age group	Number of primary vaccine doses	Number of booster doses	Interval between 1st and 2nd dose	Interval between 2nd and 3rd dose	Interval between 3rd and 4th dose
Pfizer-BioNTech	5–11 years	3	NA	3 weeks	≥4 weeks	N/A
Pfizer-BioNTech	≥12 years	3	1	3 weeks	≥4 weeks	≥3 months
Moderna	≥18 years	3	1	4 weeks	≥4 weeks	≥3 months
Janssen	≥18 years	1 Janssen, followed by 1 mRNA	1	4 weeks	≥2 months	N/A

Moderate and severe immunocompromising conditions and treatments include but are not limited to:

- Active treatment for solid tumor and hematologic malignancies
- Receipt of solid-organ transplant and taking immunosuppressive therapy
- Receipt of chimeric antigen receptor (CAR)-T-cell therapy or hematopoietic cell transplant (HCT) (within 2 years of transplantation or taking immunosuppression therapy)
- Moderate or severe primary immunodeficiency (e.g., DiGeorge syndrome, Wiskott-Aldrich syndrome)
- Advanced or untreated HIV infection (people with HIV and CD4 cell counts $<200/\text{mm}^3$, history of an AIDS-defining illness without immune reconstitution, or clinical manifestations of symptomatic HIV)
- Active treatment with high-dose corticosteroids (i.e., ≥ 20 mg prednisone or equivalent per day when administered for ≥ 2 weeks), alkylating agents, antimetabolites, transplant-related immunosuppressive drugs, cancer chemotherapeutic agents classified as severely immunosuppressive, tumor necrosis factor (TNF) blockers, and other biologic agents that are immunosuppressive or immunomodulatory